



Reducing Solar Stress on Avocados

Heat stress and sunburn caused from excessive temperatures and ultraviolet (UV) light can damage an avocado crop; significantly reducing marketable yield and cutting deep into a grower's profit. First, with grove establishment, Purshade® can protect newly planted trees from extreme conditions and accelerate the productivity of young groves. Secondly, when established trees experience extreme heat stress, photosynthetic activity can shut down, limiting overall productivity. And thirdly, direct sunburn damage on fruit, lowers yields and reduces economic returns at harvest.



Sun-damaged avocados

Purshade reduces solar stress in crops by protecting the foliage and fruit from damaging ultraviolet (UV) and infrared (IR) radiation while still allowing photosynthesis to occur. Engineered with advanced reflectance technology and based on calcium carbonate, a highly reflective mineral, Purshade has been shown to reduce sunburn damage and minimize overall heat stress. Purshade effectively reflects harmful UV and IR radiation away from plants.



Designed with the grower in mind, Purshade comes in an easy-to-handle liquid formulation that mixes quickly, applies evenly, is nonabrasive to equipment, and can be removed as required during standard post-harvest processing. The product can be applied with standard spray equipment and in most cases can be incorporated into an existing spray program.* Purshade helps maximize the value of every treated acre by reducing solar-related losses, thus increasing marketable yield, enhancing crop quality, and improving water use efficiency.


Benefits

- Assists in establishment of young trees
- Reduces culls caused by sun damage
- Improves overall pack out
- Enhances water use efficiency

Features

- Reduces damage caused by UV and IR radiation
- Easy-to-use, nonabrasive liquid formulation
- Lowers plant surface temperatures, therefore reduces plant stress and promotes biomass development
- Promotes efficient photosynthesis and does not block stomata
- Tank mixes easily, covers evenly, and can be applied with standard sprayer equipment
- Washes off during standard post-harvest processing

 AVOCADOS
SUN DAMAGE EVALUATION



Source: Agroasa, Dominican Republic
Semil Avocados. Purshade applied at 8 L/200 liters of water (4% spray concentration) to 50 trees. Fruit evaluated from 5 trees treated and 5 trees untreated. Application dates: 6/7 & 7/7/2009.

* The compatibility of Purshade with all potential tank mix partners is not knowable. Conduct a jar test first to determine compatibility and consult the product labels of the tank mix partners.

How Does Purshade Work?

Purshade plant protectant is available in a flowable suspension concentrate that is mixed with water and then sprayed directly on plant surfaces. Once dry, Purshade forms an even film of millions of microscopic “prisms” or mirrors that reflect harmful ultraviolet radiation (UV) and infrared radiation (IR) while not blocking leaf stomata, therefore not impeding photosynthesis. The reflective properties of Purshade protect fruit from direct sunburn damage and help prevent heat stress in the entire crop canopy. Keeping plants cooler, while ambient temperatures are extreme, reduces stress and enables the crop to maintain its normal photosynthetic rate longer. When Purshade is used during periods of high light and temperature extremes, crops have the solar protection needed to better reach their full potential and use available water resources more efficiently.



Application Guidelines and Rates

Purshade’s liquid formula is designed to mix easily in the tank. The product can be mixed with crop protection products (jar test recommended) and can be applied with typical ground or aerial sprayers using standard nozzles. Standard post-harvest cleaning and washing processes are generally sufficient to remove the Purshade product from the surfaces of the fruit.

APPLICATION GUIDELINES FOR AVOCADOS

Use	Gallons/Acre	Instructions
The first application of Purshade should occur prior to fruit size reaching 3/4–1 inch in diameter, or prior to extreme solar conditions.	2	Apply <u>before</u> extreme heat conditions occur.
Apply Purshade 2–3 additional times throughout the season, treating the trees approximately every 14–21 days.	2	Apply as needed based on growth and weather conditions.

Important: Always read and follow label instructions when using this product.

Use Tips

- Use the amount of water volume per acre needed to achieve uniform spray deposition. Avoid runoff.
- Use fine nozzles and the appropriate tractor speed and pressure to achieve very fine droplets that will not coalesce into larger droplets.
- Apply each application in the opposite direction of the previous application.
- Purshade is physically compatible with most crop protection products. If compatibility is unknown, perform a jar test before mixing. It is not possible to test every product or combination of products used with Purshade. If the effectiveness of a tank mixture is in doubt, apply the materials separately. Do not tank mix with products requiring a neutral or acidic pH, as the alkaline pH of Purshade might speed degradation of the other products.
- Purshade may be applied in a band spray. Use the recommended rate per acre of Purshade in a quantity of water per acre to achieve complete coverage. Adjust the band width and the volume of water based on plant size to achieve uniform coverage.
- Crops that are to be marketed fresh but have a white film of Purshade remaining at harvest may be washed. Purshade is normally removed with common washing techniques. If the crop is field packed and will not be washed, sprays should be reduced or discontinued in ample time before harvest to allow normal attrition of the film from wind, rain, and plant growth. Consult local experts for more information. Note: When Purshade applications are discontinued, the crop will begin to lose its protective coating and sunburn protection will be lost.



Sample of ideal coverage



TESSENDERLO KERLEY, INC.
2255 North 44th Street
Suite 300
Phoenix, AZ 85008-3279

Customer Service:
1-800-525-2803
1-602-889-8300

www.novasource.com
www.purshade.com

WARRANTY AND LIMITATION OF DAMAGES— Tessenderlo Kerley, Inc. warrants only that this product conforms to the product description on the label. Except as warranted by this label, Tessenderlo Kerley, Inc. makes no representation or warranty or guarantee, whether expressed or implied, of fitness for a particular purpose of merchantability, or of product performance. Tessenderlo Kerley, Inc. does not authorize any agent or representative to make any such representation, warranty or guarantee. To the extent consistent with applicable law, Tessenderlo Kerley, Inc.’s maximum liability for breach of its warranty or for use of this product, regardless of the form of action, shall be limited to the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and disposal liability resulting from handling, storage, use and disposal of this product. If buyer does not agree with or accept these warranty and liability limitations, buyer may return the unopened container to the place of purchase for full refund. Buyer’s use of this product shall constitute conclusive evidence of buyer’s acknowledgement and acceptance of the foregoing limitations. Some states do not allow the exclusion of implied warranties or the limitation of certain damages, so the above may not apply. The purchase, delivery, acceptance and use of this product by the buyer are subject to the terms and conditions of seller’s sales invoice for this product. Copyright©2012 Tessenderlo Kerley, Inc. All rights reserved. U.S. and international patents pending. Purshade and NovaSource are registered trademarks of Tessenderlo Kerley, Inc. The use of Purshade® in agricultural crop protection applications is covered by U.S. Patents 6,027,740, 6,110,867, and 6,464,995.