

Surround[®]

Agricultural Crop Protectant



Treated Plant: Bananas planted in October 2011 and treated with SURROUND every 8 days in standard sigatoka program. (11 applications)



Untreated Control: Both photos taken 2/23/13 at similar distance and angle.

Sunlight is essential for banana growth and fruit production, but excessive exposure to solar radiation can result in heat stress to both young banana plants during establishment and to mature plants during the lifetime of this perennial crop. The highly engineered kaolin particles in **Surround[®] WP Crop Protectant** reflect infrared and ultraviolet radiation, significantly reducing heat stress and excessive light damage. A visual symptom of adverse plant stress is banana leaves hanging limply from the central leaf vein in a wilted condition. By reflecting excessive radiation, SURROUND maintains plants in more optimum and productive growing conditions.

CORBANA Test Results

The multi-year, multi-generation CORBANA trial was conducted in Costa Rica under heavy rainfall, water shortage conditions and high and low temperatures. In order to minimize the effect of stress on cv Grande Naine banana plants, SURROUND behavior on productivity and quality was evaluated at rates of 3% and 5% every 8 days, and 5% every 15 days compared to a control treatment in mixtures with fungicides for black Sigatoka (*Mycosphaerella fijiensis*) control. Vegetative and productivity variables were evaluated for four generations at the CORBANA Research Center from 2011 to 2014.

Results showed greater vegetative vigor and a lower interval of days between planting to flowering (12 to 15 fewer days) in banana plants applied with SURROUND, and higher return rates (number of fruit per hectare per year) – even under conditions of water stress and low and high temperatures where observed. Treatments with SURROUND had greater numbers of hands per bunch (0.1 to 0.6 more hands) and an increase in fruit weight (plus 1.1 to 2.0 kg/bunch) with respect to the control treatment. The fruit yield index or “ratio” was higher (plus 0.15) in plants applied with SURROUND compared to the control treatment. Evaluation of the bunch profile showed higher net weight of the fruits in the treatments applied with SURROUND (plus 0.4 to 3.4 kg/bunch).

SURROUND effectively reduced stress conditions in banana plants under adverse weather conditions such as drought and high and low temperature, translating to significantly increased productivity.

Results from other on-farm demonstration trials further validate the benefits of using SURROUND for banana plant establishment and overall productivity. In another trial conducted in 2012 in Costa Rica, SURROUND applied at 5% w/v every 15 days (Table 2) produced increased vegetative growth in mother and daughter plants at the time of shooting.

Table 1. SURROUND increased average productivity During all generations evaluated R0 to R3. (CORBANA, Costa Rica, 2011-2014)

Variable	Surround WP	Control
Number of Hands/ Bunch	7	6.65
Bunch Weight (kg)	26.29	24.8
Yield Ratio (boxes/bunch)	1.25	1.16
Days Avg. from Planting to Flowering	245	239
Cum. Avg. Days from Planting to Flowering	13	

Note: Data in Table 1 represents average data for all generations at different rates and intervals.

Table 2. Surround increased vegetative growth in mother and daughter plants at the time of shooting (Bioagro, DeLeyte Farm, Costa Rica, 2012. Surround at 5% w/v every 15 days)

Variable	Surround WP	Control
Number of Leaves – Mother Plant	13.05a	12.6b
Plant Height – Daughter Plant (cm)	153.5a	137.57b
Number of Leaves – Daughter Plant	9.64a	8.32b
Days from Transplant to Shooting	171a	177b

*Means followed by different letters are significantly different (p<0.05).

TESSENDERLO KERLEY, INC.
2255 North 44th Street
Suite 300
Phoenix, AZ 85008-3279

Customer Service:
800-525-2803
602-889-8300

novasource.com

The Science of Reflecting Light and Heat

SURROUND was co-developed with scientists at the U.S. Department of Agriculture (USDA). It is registered in the United States by the Environmental Protection Agency (EPA) and kaolin is Annex I listed in the European Union by the European Food Safety Authority (EFSA). Therefore, use of SURROUND for export markets is free of tolerance complications. Also, SURROUND is listed for use in organic crops by the Organic Materials Review Institute (OMRI). The patented kaolin particles in SURROUND reflect excessive infrared and ultraviolet radiation from the plant canopy, and do so more efficiently than other solar protectant products. A study evaluating the solar reflectance capabilities of various products shows SURROUND is superior at reflecting the most damaging wavelengths of solar radiation compared to the other products tested. For more information, visit novasource.com.

Guidelines for Use

- **Concentration:** Use 5% weight/volume or up to 10% depending on spray interval in typical spray volume for banana crop protection. For wider application intervals higher concentrations are recommended.
- **Timing:** Initiate applications for heat stress at transplanting, and continue weekly or biweekly thereafter in the normal Sigatoka disease control and crop protection program.
- **Mode of Application:** Back pack sprayer, motorized sprayer, or aircraft aerial sprayer. Calibrate sprayer and spray according to standard banana spray program volume.
- **Tank Mixing:** SURROUND may be mixed and applied with fungicides, insecticides, and foliar fertilizers because it is chemically and toxicologically inert. As a precaution, prior to tank mixing, conduct a small jar test for signs of physical incompatibility. Always add SURROUND to the tank first, and then add the other product(s) to be used in the application. Maintain agitation and mix thoroughly.
- **Post Harvest:** After harvest but before bananas are wet, ensure there are no white residues by visually inspecting bunches. Trace residues can be removed by gently wiping with a cloth.



Important: Always read the label before buying and follow label instructions when using this product.

The use of SURROUND in agricultural crop protection applications is covered by patents held by Tessenderlo Kerley Inc. Visit our website for current patent information: novasource.com.

CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Tessenderlo Kerley, Inc. makes no representation or warranty or guarantee, whether expressed or implied, of fitness for a particular purpose of merchantability, or of product performance. Tessenderlo Kerley, Inc. does not authorize any agent or representative to make any such representation, warranty or guarantee. To the extent consistent with applicable law, Tessenderlo Kerley, Inc.'s maximum liability for breach of its warranty or for use of this product, regardless of the form of action, shall be limited to the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and disposal liability resulting from handling, storage, use and disposal of this product. If buyer does not agree with or accept these warranty and liability limitations, buyer may return the unopened container to the place of purchase for full refund. Buyer's use of this product shall constitute conclusive evidence of buyer's acknowledgement and acceptance of the forgoing limitations. Some states do not allow the exclusion of implied warranties or the limitation of certain damages, so the above may not apply. The purchase, delivery, acceptance and use of this product by the buyer are subject to the terms and conditions of seller's sales invoice for this product.



TESSENDERLO KERLEY, INC.
2255 North 44th Street
Suite 300
Phoenix, AZ 85008-3279

Customer Service:
800-525-2803
602-889-8300

novasource.com