

USE GROUP 7 LINEX® FOR WATERHEMP CONTROL IN SOYBEANS

Linex®
Agricultural Herbicide

An Effective Tool for Controlling Waterhemp in Soybeans

Waterhemp is one of the most devastating weeds affecting soybean yields in the U.S. today.

This broadleaf summer annual emerges throughout the growing season, outlasting many pre-emergent herbicides.

Even non-residual post-emergent herbicides may miss late breaking weeds.

When an infestation reaches or exceeds 20 weeds per square foot, especially late in the season, the consequences can be catastrophic — reducing yields by up to 44 percent. Waterhemp that emerges as late as the V5 stage can in soybeans can cut yields by up to 10 percent.¹

Waterhemp has demonstrated resistance to six different herbicide groups, including: Group 2, Group 4, Group 5,

Group 9, Group 14 and Group 27. And resistance to multiple herbicide groups in some plant populations is now occurring in many states.

Waterhemp grows quickly and competes for sunlight and nutrients while producing an unusually high number of seeds per plant — from 250,000 to 1 million seeds under optimal conditions.²

Linex® 4L — An Alternative Mode of Action with Late-season Success for Control of Waterhemp

- Linex 4L is a Group 7 herbicide.
- LINEX 4L is effective on waterhemp and other important weeds, where rotating chemistry is an effective way to combat resistance.
- LINEX 4L is less soluble than many herbicides, which enables it to control weeds, like waterhemp, that emerge later in the season.
- The insolubility of LINEX 4L enables it to retain effectiveness in areas of heavy rainfall.
- LINEX 4L is an effective pre-emergent tool in the battle to control waterhemp. Pre-emergence programs have been found to be more effective in maximizing yields and return on investment than post-emergence programs.
- University and consultant studies have shown LINEX 4L to provide waterhemp control ranging between fair and excellent, depending upon soil type, rate of application and weed pressure.
- As the percentage of organic matter in the soil increases, the rate of LINEX 4L applications needs to increase to maintain efficacy. Do not exceed 2 pints per acre of LINEX 4L in any application.

¹ Waterhemp Management in Soybeans," available at TakeActionOnWeeds.com

² Waterhemp Management in Soybeans," available at TakeActionOnWeeds.com



Waterhemp



TESSENDERLO KERLEY, INC.

2910 North 44th Street
Suite 100
Phoenix, AZ 85018-7272

CUSTOMER SERVICE:

800-525-2803
602-889-8300

novasource.com

USE GROUP 7 LINEX® FOR WATERHEMP CONTROL IN SOYBEANS

Linex®
Agricultural Herbicide

PRE Followed by POST on Glyphosate-tolerant Soybeans		
Situation	Recommendation	
	Preemergence	Postemergence
To provide added residual activity or burndown activity on emerged weeds, preemergence applications of LINEX 4L may be tank mixed with full or reduced rates of other products registered for use in soybeans.	LINEX 4L 1.0-2.0 pint / Acre Rate per acre depends upon: <ul style="list-style-type: none">• Soil texture and organic matter• Level of residual control desired	Glyphosate-tolerant soybean varieties. Your favorite glyphosate-based POST program.
Application Information		
<ul style="list-style-type: none"> • LINEX 4L must be applied before the soybean crop emerges. • When using LINEX 4L for burndown, always add crop oil concentrate at 1 percent v/v (1 gallon per 100 gallons of spray solution) and 0.375 ounce per acre Synchrony® XP or 1 pint per acre 2,4-D LVE. 2,4-D applications must be made at least 7 days prior to planting. • For burndown of weeds not controlled by LINEX 4L, other burndown products may be tank mixed with LINEX 4L. 		



Weeds Controlled Pre-emergence		
Broadleaf Weeds	Grasses	LINEX 4L will provide partial control (suppression) of the following hard to control weeds
Amaranth, Palmer (<i>Amaranthus palmeri</i>) Carpetweed (<i>Mollugo verticillata</i>) Chickweed, Common (<i>Stellaria media</i>) Dog Fennel (<i>Mayweed Chamomile</i>) (<i>Anthemis cotula</i>) Galinsoga (<i>Galinsoga sp.</i>) Nettleleaf Goosefoot (<i>Chenopodium murale</i>) Lambsquarters (<i>Chenopodium album</i>) Mustard, Tumble (<i>Sisymbrium altissimum</i>) Mustard, Blue (<i>Chorispora tenella</i>) Mustard, Wallflower (Treacle) (<i>Erysimum cheiranthoides</i>) Pigweed, Prostrate (<i>Amaranthus blitoides</i>) Pigweed, Redroot (<i>Amaranthus retroflexus</i>) Pigweed, Smooth (<i>Amaranthus hybridus</i>) Prickley Lettuce (<i>Lactuca serriola</i>) Purslane, Common (<i>Portulaca oleracea</i>) Radish, Wild (<i>Raphanus raphanistrum</i>) Ragweed, Common (<i>Ambrosia artemisiifolia</i>) Shepherd's-purse (<i>Capsella bursa-pastoris</i>) Smartweed, Pennsylvania (<i>Polygonum pennsylvanicum</i>) Tansymustard, Pinnate (<i>Descurainia pinnata</i>) Waterhemp, Common (<i>Amaranthus rudis</i>) Waterhemp (<i>Amaranthus tuberculatus</i>)	Barnyardgrass (Watergrass) [<i>Echinochloa crus-galli</i>] Canarygrass (<i>Phalaris canariensis</i>) Large Crabgrass (<i>Digitaria sanguinalis</i>) Green Foxtail (<i>Setaria viridis</i>) Yellow Foxtail (<i>Setaria glauca</i>) Goosegrass (<i>Eleusine indica</i>) Fall Panicum (<i>Panicum dichotomiflorum</i>)	Annual Morningglories only See <i>Ipomoae spp.</i> Entire Leaf Morningglory (<i>Ipomoae hederacea integriflora</i>) Ivy Leaf Morningglory (<i>Ipomoae hederacea</i>) Pitted Morningglory (<i>Ipomoae ipola & I. lacunosa</i>) Cocklebur, Common (<i>Xanthium strumarium</i>) Dayflower, Common (Asiatic) (<i>Commelinia communis</i>) Eastern Black Nightshade (<i>Solanum ptychanthrum</i>) Florida Beggarweed (<i>Desmodium tortuosum</i>) Florida Pursley (Florida Purslane) (<i>Richardia scabra</i>) Giant Foxtail (<i>Setaria faberii</i>) Prickly Sida (<i>Sida spinosa</i>) Ragweed, Giant (<i>Ambrosia trifida</i>) Sicklepod Cassia (<i>obtusifolia</i>) Smooth Crabgrass (<i>Digitaria ischaemum</i>) Velvetleaf (Buttonweed) (<i>Abutilon theophrasti</i>) Waterhemp, Common (<i>Amaranthus rudis</i>)

Important: Always read the label before buying and follow label instructions when using this product.

For more information, visit novasource.com.

WARRANTY AND LIMITATION OF DAMAGES

Tessenderlo Kerley, Inc. (TKI) warrants only that this product conforms to the product description on the label. Except as warranted by this label, TKI makes no representation or warranty or guarantee, whether expressed or implied, of fitness for a particular purpose of merchantability, or of product performance. TKI does not authorize any agent or representative to make any such representation, warranty or guarantee. To the extent consistent with applicable law, TKI's maximum liability for breach of its warranty or for use of this product, regardless of the form of action, shall be limited to the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and disposal liability resulting from handling, storage, use and disposal of this product. If buyer does not agree with or accept these warranty and liability limitations, buyer may return the unopened container to the place of purchase for full refund. Buyer's use of this product shall constitute conclusive evidence of buyer's acknowledgement and acceptance of the foregoing limitations. Some jurisdictions do not allow the exclusion of implied warranties or the limitation of certain damages, so the above may not apply. The purchase, delivery, acceptance and use of this product by the buyer are subject to the terms and conditions of seller's sales invoice for this product.

NovaSource® and Linex® are registered trademarks of Tessenderlo Kerley, Inc.

All other brands and trade names listed are registered trademarks of their respective companies.

© 2006, 2009, 2016, 2021 Tessenderlo Kerley, Inc. All rights reserved.

2022-01-04T-Linex-PI-Waterhemp

NovaSource®
Tessenderlo Group

TESSENDERLO KERLEY, INC.

2910 North 44th Street

Suite 100

Phoenix, AZ 85018-7272

CUSTOMER SERVICE:

800-525-2803

602-889-8300

novasource.com